


		CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT FOR INVOICE 12345 DATED 01/01/17				
J R SMITH CALF GROWER CATTLE FEED MEDICATED TYPE B FEED CONTAINING AMPROLIUM						
FORMULA #: JRSMITHJAN2017						
STATEMENT OF PURPOSE						
As an aid in treatment of coccidiosis caused by Eimeria bovis and E. zuernii in calves.						
ACTIVE DRUG INGREDIENT						
Amprolium		5113.64 grams/ton 2556.82 milligrams/lb. 0.56 %				
USE DIRECTIONS						
Use 10 pounds of 2.5% Type B per ton of feed to produce a Type C medicated feed containing 0.0125% amprolium. Use 1,000 pounds of 2.5% Type B per ton to produce a Type C medicated feed containing 1.25% amprolium.						
<i>Feed the resulting Type C medicated feed in amounts according to the chart below to provide 454mg amprolium/100 pounds of bodyweight per day for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely the rest of the group has been exposed and all calves in the group should be treated.</i>						
Calf Weight (pounds)	amprolium required(mg)	lbs. to feed per day		Calf Weight (pounds)	amprolium required(mg)	lbs. to feed per day
100	454	0.18		300	1362	0.53
150	681	0.27		350	1589	0.62
200	908	0.36		400	1816	0.71
250	1135	0.44		450	2043	0.80
NOTE: For satisfactory diagnosis a microscopic examination of the feces should be done by a veterinarian or diagnostic laboratory before treatment. When treating outbreaks, the drug should be administered promptly after diagnosis is determined.						
PRECAUTIONARY STATEMENTS						
WARNING: Withdraw 24 hours before slaughter. Use as sole source of Amprolium. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.						
CAUTION: Do not use Amprolium in feeds containing bentonite.						
MANUFACTURED BY: BLUEBIRD FEED MILL, ANY CITY ANY STATE, 55555						
Net Weight on Bag and/or Invoice						
Revised 12/01/16						

		CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT FOR INVOICE 12345 DATED 01/01/17				
J R SMITH CALF GROWER CATTLE FEED MEDICATED TYPE B FEED CONTAINING AMPROLIUM						
FORMULA #: JRSMITHJAN2017						
STATEMENT OF PURPOSE						
As an aid in treatment of coccidiosis caused by Eimeria bovis and E. zuernii in calves.						
ACTIVE DRUG INGREDIENT						
Amprolium		5113.64 grams/ton 2556.82 milligrams/lb. 0.56 %				
USE DIRECTIONS						
Use 10 pounds of 2.5% Type B per ton of feed to produce a Type C medicated feed containing 0.0125% amprolium. Use 1,000 pounds of 2.5% Type B per ton to produce a Type C medicated feed containing 1.25% amprolium.						
<i>Feed the resulting Type C medicated feed in amounts according to the chart below to provide 454mg amprolium/100 pounds of bodyweight per day for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely the rest of the group has been exposed and all calves in the group should be treated.</i>						
Calf Weight (pounds)	amprolium required(mg)	lbs. to feed per day		Calf Weight (pounds)	amprolium required(mg)	lbs. to feed per day
100	454	0.18		300	1362	0.53
150	681	0.27		350	1589	0.62
200	908	0.36		400	1816	0.71
250	1135	0.44		450	2043	0.80
NOTE: For satisfactory diagnosis a microscopic examination of the feces should be done by a veterinarian or diagnostic laboratory before treatment. When treating outbreaks, the drug should be administered promptly after diagnosis is determined.						
PRECAUTIONARY STATEMENTS						
WARNING: Withdraw 24 hours before slaughter. Use as sole source of Amprolium. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.						
CAUTION: Do not use Amprolium in feeds containing bentonite.						
MANUFACTURED BY: BLUEBIRD FEED MILL, ANY CITY ANY STATE, 55555						
Net Weight on Bag and/or Invoice						
Revised 12/01/16						