


	<b>CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT</b> FOR INVOICE 50000 DATED 01/01/17	
<b>JAMES SMITH GROWER PHEASANT FEED</b> <b>MEDICATED</b> TYPE C FEED CONTAINING AMPROLIUM  FORMULA #: SMITHPHEASANTJAN17		
<b>STATEMENT OF PURPOSE</b>		
For the prevention of coccidiosis in growing pheasants caused by Eimeria colchici, E. duodenalis, and E. phasiani		
<b>ACTIVE DRUG INGREDIENT</b>		
Amprolium 159.09 grams/ton 79.55 milligrams/pound 0.0175 %		
<b>USE DIRECTIONS</b>		
Feed continuously as sole ration.		
<b>PRECAUTIONARY STATEMENTS</b>		
<b>WARNING:</b> Use as the sole source of amprolium.  Do not change the litter while giving this feed unless absolutely necessary.  If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis, and follow the instructions of your veterinarian or poultry pathologist.  Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.  Fertility, hatchability, and other reproductive data are not available on amprolium in breeding pheasants.  <b>CAUTION:</b> Do not use in feeds containing bentonite.		
MANUFACTURED BY:  BLUE BIRD FEED MILL, ANY CITY, ANY STATE, 55555  Net Weight on Bag and/or Invoice		
Label Revised 12/01/16		

	<b>CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT</b> FOR INVOICE 50000 DATED 01/01/17	
<b>JAMES SMITH PHEASANT FEED</b> <b>MEDICATED</b> TYPE C FEED CONTAINING AMPROLIUM  FORMULA #: SMITHPHEASANTJAN17		
<b>STATEMENT OF PURPOSE</b>		
For the prevention of coccidiosis caused by Eimeria colchici, E. duodenalis, and E. phasiani		
<b>ACTIVE DRUG INGREDIENT</b>		
Amprolium 159.09 grams/ton 79.55 milligrams/pound 0.0175 %		
<b>USE DIRECTIONS</b>		
Feed continuously as sole ration.		
<b>PRECAUTIONARY STATEMENTS</b>		
<b>WARNING:</b> Use as the sole source of amprolium.  Do not change the litter while giving this feed unless absolutely necessary.  If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis, and follow the instructions of your veterinarian or poultry pathologist.  Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.  Fertility, hatchability, and other reproductive data are not available on amprolium in breeding pheasants.  <b>CAUTION:</b> Do not use in feeds containing bentonite.		
MANUFACTURED BY:  BLUE BIRD FEED MILL, ANY CITY, ANY STATE, 55555  Net Weight on Bag and/or Invoice		
Label Revised 12/01/16		