


	CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT FOR INVOICE 50000 DATED 01/01/17							
JAMES SMITH LAYER CHICKEN FEED MEDICATED TYPE C FEED CONTAINING AMPROLIUM FORMULA #: SMITHLAYERJAN17								
STATEMENT OF PURPOSE								
For development of active immunity to coccidiosis in replacement chickens up to 5 weeks of age under conditions of severe exposure to coccidiosis. For the prevention and treatment of moderate outbreaks of coccidiosis in laying chickens.								
ACTIVE DRUG INGREDIENT								
<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 60%;">Amprolium</td> <td style="text-align: right;">113.52 grams/ton</td> </tr> <tr> <td></td> <td style="text-align: right;">56.76 milligrams/pound</td> </tr> <tr> <td></td> <td style="text-align: right;">0.0125 %</td> </tr> </table>			Amprolium	113.52 grams/ton		56.76 milligrams/pound		0.0125 %
Amprolium	113.52 grams/ton							
	56.76 milligrams/pound							
	0.0125 %							
USE DIRECTIONS								
<p>For the development of active immunity in replacement chickens: Feed continuously until onset of production.</p> <p>Prevention of cocci in laying hens: Use 0.0125% amprolium Type C medicated feed continuously. Treatment of moderate cocci outbreak in laying hens: Feed 0.0125% amprolium Type C medicated feed for two weeks. Treatment of severe outbreaks in laying hens: Feed 0.025% amprolium Type C medicated for one week.</p>								
PRECAUTIONARY STATEMENTS								
<p>WARNING: Use as the sole source of amprolium.</p> <p>Do not change the litter while giving this feed unless absolutely necessary.</p> <p>If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis, and follow the instructions of your veterinarian or poultry pathologist.</p> <p>Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.</p> <p>In replacement flocks the grower must expect that excessive exposure to one or more species may overwhelm the drug in some flocks and prompt treatment will be required.</p> <p>CAUTION: Do not use with feeds containing bentonite.</p>								
MANUFACTURED BY: BLUE BIRD FEED MILL, ANY CITY, ANY STATE, 55555 Net Weight on Bag and/or Invoice								
Label Revised 07/26/17								

	CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT FOR INVOICE 50000 DATED 01/01/17							
JAMES SMITH LAYER CHICKEN FEED MEDICATED TYPE C FEED CONTAINING AMPROLIUM FORMULA #: SMITHLAYERJAN17								
STATEMENT OF PURPOSE								
For development of active immunity to coccidiosis in replacement chickens up to 5 weeks of age under conditions of severe exposure to coccidiosis. For the prevention and treatment of moderate outbreaks of coccidiosis in laying chickens.								
ACTIVE DRUG INGREDIENT								
<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 60%;">Amprolium</td> <td style="text-align: right;">113.52 grams/ton</td> </tr> <tr> <td></td> <td style="text-align: right;">56.76 milligrams/pound</td> </tr> <tr> <td></td> <td style="text-align: right;">0.0125 %</td> </tr> </table>			Amprolium	113.52 grams/ton		56.76 milligrams/pound		0.0125 %
Amprolium	113.52 grams/ton							
	56.76 milligrams/pound							
	0.0125 %							
USE DIRECTIONS								
<p>For the development of active immunity in replacement chickens: Feed continuously until onset of production.</p> <p>Prevention of cocci in laying hens: Use 0.0125% amprolium Type C medicated feed continuously. Treatment of moderate cocci outbreak in laying hens: Feed 0.0125% amprolium Type C medicated feed for two weeks. Treatment of severe outbreaks in laying hens: Feed 0.025% amprolium Type C medicated for one week.</p>								
PRECAUTIONARY STATEMENTS								
<p>WARNING: Use as the sole source of amprolium.</p> <p>Do not change the litter while giving this feed unless absolutely necessary.</p> <p>If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis, and follow the instructions of your veterinarian or poultry pathologist.</p> <p>Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.</p> <p>In replacement flocks the grower must expect that excessive exposure to one or more species may overwhelm the drug in some flocks and prompt treatment will be required.</p> <p>CAUTION: Do not use with feeds containing bentonite.</p>								
MANUFACTURED BY: BLUE BIRD FEED MILL, ANY CITY, ANY STATE, 55555 Net Weight on Bag and/or Invoice								
Label Revised 07/26/17								