CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT		
FOR INVOICE 90101 DATED 01/01/17		
FOR INVOICE 90101 DATED 01/01/17		
JAMES SMITH DAIRY HEIFER GROWER CATTLE FEED OBSERVE LABEL DIRECTIONS		
MEDICATED		
SUPPLEMENT FEED CONTAINING THE DRUGS CHLORTETRACYCLINE AND DECOQUINATE		
CAUTION: Federal law restricts medicated feed containing this VFD drug to use by or on the order of a		
licensed veterinarian.		
VFD Expiration Date: Saturday, January 7, 2017		
Formula # JS DHG Dec 16		
STATEMENT OF PURPOSE		
For prevention of coccidiosis caused by Eimeria bovis and Eimeria zuernii. For		
treatment of bacterial enteritis caused by Escherichia coli and for treatment of		
bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to		
Chlortetracycline.		
ACTIVE DRUG INGREDIENT		
Chlortetracycline (as chlortetracycline calcium complex) 1000.00 grams/ton		
500.00 mg/pound		
Decoquinate		
11.35 mg/pound		
USE DIRECTIONS		
Feed supplement as a top dress or mix into daily ration to provide 22.7 mg decoquinate and 1 g		
chlortetracycline per 100 lb. bodyweight per day for not more than 5 days.		
<b>2.00</b> pounds of this supplement will medicate 100 pounds of body weight.		
NOTICE: This feed may not be fed after Saturday, January 7, 2017		
PRECAUTIONARY STATEMENTS		
WARNING: Do not feed to animals producing milk for food. A withdrawal period has not been established		
for this product in pre-ruminating calves. Do not feed to calves to be processed for veal. Withdraw 24 hours		
prior to slaughter when manufactured from ChlorMax Type A medicated articles and Pennchlor Type A		
medicated articles. Zero withdrawal time when manufactured from Aureomycin Type A medicated articles		
manufactured by Zoetis.		
<b>CAUTION:</b> Do not use decoquinate in feeds containing bentonite.		
LIMITATIONS FOR USE: Feed Type C feed at a rate to provide 22.7 mg decoguinate and 1 g		
chlortetracycline per 100 lb. of bodyweight (0.5 mg/kg) per day for not more than 5 days. When consumed,		
feed 22.7 mg decoquinate per 100 lb. bodyweight per day for total of 28 days to prevent coccidiosis.		
MANUFACTURED BY:		
BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555		
Net Weight on Bag and/or Invoice		
Net Weight on bay and/or invoice		

revision 12/01/16

CUSTOMER-FORMULA MEDICATE FOR INVOICE 90101	ED LABEL ATTACHMENT DATED 01/01/17	
FOR INVOICE 90101	DATED UNUNTY	
JAMES SMITH DAIRY HEIFER GROWER MEDICATE	DIDECTIONS	
SUPPLEMENT FEED CONTAINING THE DRUGS CHLORTETRACYCLINE AND DECOQUINATE		
CAUTION: Federal law restricts medicated feed containing this VFD drug to use by or on the order of a licensed veterinarian.		
VFD Expiration Date: Saturday, January 7, 2017		
Formula # JS DHG Dec 16		
STATEMENT OF P		
For prevention of coccidiosis caused by Eimeria bovis and Eimeria zuernii. For		
treatment of bacterial enteritis caused by Escherichia coli and for treatment of		
bacterial pneumonia caused by <i>Pasteurella mu</i> Chlortetracycline.	ultocida organisms susceptible to	
ACTIVE DRUG INGREDIENT		
Chlortetracycline (as chlortetracycline calcium complex)	1000.00 grams/ton	
	500.00 mg/pound	
Decoquinate	22.70 grams/ton	
	11.35 mg/pound	
USE DIRECTIONS		
Feed supplement as a top dress or mix into daily ration to provide 22.7 mg decoquinate and 1 g		
chlortetracycline per 100 lb. bodyweight per day for not more than 5 days.		
2.00 pounds of this supplement will medicate 10	0 pounds of body weight.	
NOTICE: This feed may not be fed after Saturday, January 7, 2017		
PRECAUTIONARY ST	ATEMENTS	
WARNING: Do not feed to animals producing milk for food. A withdrawal period has not been established		
for this product in pre-ruminating calves. Do not feed to calves to be processed for veal. Withdraw 24 hours		
prior to slaughter when manufactured from ChlorMax Type A medicated articles and Pennchlor Type A		
medicated articles. Zero withdrawal time when manufactured from Aureomycin Type A medicated articles		
manufactured by Zoetis.		
CAUTION: Do not use decoquinate in feeds containing bentonite.		
LIMITATIONS FOR USE: Feed Type C feed at a rate to pr		
chlortetracycline per 100 lb. of bodyweight (0.5 mg/kg) per day for not more than 5 days. When consumed, feed 22.7 mg decoquinate per 100 lb. bodyweight per day for total of 28 days to prevent coccidiosis.		
MANUEACTIO		
MANUFACTURED BY: BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555		
Net Weight on Bag and/or Invoice		
Net Weight on Bag and/or involce		

revision 12/01/16