



<p>CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT FOR INVOICE 90101 DATED 01/01/17</p> <p>Blue Sky Dairy Heifer Grower CATTLE FEED MEDICATED</p> <p>COMPLETE FEED CONTAINING THE DRUG DECOQUINATE AND THE FEED ADDITIVE DIFLUBENZURON</p> <p>Formula # BSDHG Nov 16</p>	<p>TAKE TIME  OBSERVE LABEL DIRECTIONS</p>
STATEMENT OF PURPOSE	
<p>For the prevention of coccidiosis in cattle (including ruminating and non-ruminating calves and veal calves) caused by <i>Eimeria bovis</i> and <i>E. zuernii</i>.</p> <p>For the prevention of adult house, stable, face and horn fly, emergence from the manure of treated cattle.</p>	
ACTIVE DRUG AND FEED ADDITIVE INGREDIENTS	
Decoquinate	13.18 grams/ton 6.59 mg/pound
Diflubenzuron	2.64 grams/ton 1.32 mg/pound
USE DIRECTIONS	
<p>Feed at a rate of 22.7 mg decoquinate per 100 lb of body weight (0.5 mg/kg) per day. Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard. The required feeding rate of diflubenzuron is 0.1 mg/kg of body weight per day.</p> <p>3.44 pounds of this feed will provide 22.7 mg decoquinate and 4.55 mg diflubenzuron per 100 pounds of body weight.</p>	
PRECAUTIONARY STATEMENTS	
<p>WARNING: Do not feed to animals producing milk for human consumption. CAUTION: Do not use in feeds containing bentonite. LIMITATIONS: Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard.</p> <p style="text-align: center;">CAUTION-KEEP OUT OF REACH OF CHILDREN</p> <p>See the attached Diflubenzuron label related to safety (ie. Personal protective equip., first aid, environmental hazards etc.).</p>	
MANUFACTURED BY:	
BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555	
Net Weight on Bag and/or Invoice	revision 12/01/16

<p>CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT FOR INVOICE 90101 DATED 01/01/17</p> <p>Blue Sky Dairy Heifer Grower CATTLE FEED MEDICATED</p> <p>COMPLETE FEED CONTAINING THE DRUG DECOQUINATE AND THE FEED ADDITIVE DIFLUBENZURON</p> <p>Formula # BSDHG Nov 16</p>	<p>TAKE TIME  OBSERVE LABEL DIRECTIONS</p>
STATEMENT OF PURPOSE	
<p>For the prevention of coccidiosis in cattle (including ruminating and non-ruminating calves and veal calves) caused by <i>Eimeria bovis</i> and <i>E. zuernii</i>.</p> <p>For the prevention of adult house, stable, face and horn fly, emergence from the manure of treated cattle.</p>	
ACTIVE DRUG AND FEED ADDITIVE INGREDIENTS	
Decoquinate	13.18 grams/ton 6.59 mg/pound
Diflubenzuron	2.64 grams/ton 1.32 mg/pound
USE DIRECTIONS	
<p>Feed at a rate of 22.7 mg decoquinate per 100 lb of body weight (0.5 mg/kg) per day. Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard. The required feeding rate of diflubenzuron is 0.1 mg/kg of body weight per day.</p> <p>3.44 pounds of this feed will provide 22.7 mg decoquinate and 4.55 mg diflubenzuron per 100 pounds of body weight.</p>	
PRECAUTIONARY STATEMENTS	
<p>WARNING: Do not feed to animals producing milk for human consumption. CAUTION: Do not use in feeds containing bentonite. LIMITATIONS: Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard.</p> <p style="text-align: center;">CAUTION-KEEP OUT OF REACH OF CHILDREN</p> <p>See the attached Diflubenzuron label related to safety (ie. Personal protective equip., first aid, environmental hazards etc.).</p>	
MANUFACTURED BY:	
BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555	
Net Weight on Bag and/or Invoice	revision 12/01/16