

CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT
FOR INVOICE 90101 DATED 01/01/17



**Blue Sky Dairy Heifer Grow Supp CATTLE FEED
MEDICATED**

FEED SUPPLEMENT CONTAINING THE DRUG DECOQUINATE AND
THE FEED ADDITIVE DIFLUBENZURON

Formula # BSDHG Nov 16

STATEMENT OF PURPOSE

For the prevention of coccidiosis in cattle (including ruminating and non-ruminating calves and veal calves) caused by *Eimeria bovis* and *E. zuernii*.

For the prevention of adult house, stable, face and horn fly, emergence from the manure of treated cattle.

ACTIVE DRUG AND FEED ADDITIVE INGREDIENTS

Decoquinate	90.91 grams/ton 45.46 mg/pound
Diflubenzuron	18.27 grams/ton 9.14 mg/pound

USE DIRECTIONS

Feed at a rate of 22.7 mg decoquinate per 100 lb of body weight (0.5 mg/kg) per day.
Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard. The required feeding rate of diflubenzuron is 0.1 mg/kg of body weight per day.

Administer as a top dress supplement or mix into the daily ration to provide 22.7 mg per pound per day.

0.5 pounds of this supplement will provide 22.7 mg decoquinate per 100 lb of bodyweight.

PRECAUTIONARY STATEMENTS

WARNING: Do not feed to animals producing milk for human consumption.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS: Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard.

CAUTION-KEEP OUT OF REACH OF CHILDREN

See the attached Diflubenzuron label related to safety (ie. Personal protective equip., first aid, environmental hazards etc.).

MANUFACTURED BY:

BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555

Net Weight on Bag and/or Invoice

revision 12/01/16

CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT
FOR INVOICE 90101 DATED 01/01/17



**Blue Sky Dairy Heifer Grow Supp CATTLE FEED
MEDICATED**

FEED SUPPLEMENT CONTAINING THE DRUG DECOQUINATE AND
THE FEED ADDITIVE DIFLUBENZURON

Formula # BSDHG Nov 16

STATEMENT OF PURPOSE

For the prevention of coccidiosis in cattle (including ruminating and non-ruminating calves and veal calves) caused by *Eimeria bovis* and *E. zuernii*.

For the prevention of adult house, stable, face and horn fly, emergence from the manure of treated cattle.

ACTIVE DRUG AND FEED ADDITIVE INGREDIENTS

Decoquinate	90.91 grams/ton 45.46 mg/pound
Diflubenzuron	18.27 grams/ton 9.14 mg/pound

USE DIRECTIONS

Feed at a rate of 22.7 mg decoquinate per 100 lb of body weight (0.5 mg/kg) per day.
Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard. The required feeding rate of diflubenzuron is 0.1 mg/kg of body weight per day.

Administer as a top dress supplement or mix into the daily ration to provide 22.7 mg per pound per day.

0.5 pounds of this supplement will provide 22.7 mg decoquinate per 100 lb of bodyweight.

PRECAUTIONARY STATEMENTS

WARNING: Do not feed to animals producing milk for human consumption.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS: Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard.

CAUTION-KEEP OUT OF REACH OF CHILDREN

See the attached Diflubenzuron label related to safety (ie. Personal protective equip., first aid, environmental hazards etc.).

MANUFACTURED BY:

BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555

Net Weight on Bag and/or Invoice

revision 12/01/16