

CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT
 FOR INVOICE 50001 DATED 1/1/2017



**Replacement Heifer CATTLE FEED
 MEDICATED**
 SUPPLEMENT FEED CONTAINING RUMENSIN™ (MONENSIN)

FORMULA # Doe RH Dec 2016

STATEMENT OF PURPOSE

For increased rate of weight gain; for the prevention and control of coccidiosis due to Eimeria bovis and E. zuernii in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

Monensin (as monensin sodium) 75.00 grams/ton
 37.50 milligrams/pound

USE DIRECTIONS

NOTE: Monensin must fed with at least 1 pound of feed. If indicated feeding rates are less than 1 pound, mix additional non-medicated feed to make at least one (1) pound of feed.

FOR INCREASED RATE OF WEIGHT GAIN:

Feed so that cattle will be consuming between 1.33 and 5.33 pounds per head per day in not less than 1lb. of feed to provide not less than 50 nor more than 200 milligrams per head per day, or after the 5th day feed at the rate of 10.67 pounds per head per every other day in not less than 2 pounds of feed to supply 400 milligrams monensin per head every other day.

IMPORTANT: For cattle consuming monensin for the first time, during the first 5 days of feeding, cattle should receive no more than 100 mg monensin or 2.67 pounds of this feed/head/day.

FOR COCCIDIOSIS PREVENTION AND CONTROL:

Hand feed this ration daily in accordance with the feeding schedule below to supply 0.14 to 0.42 mg monensin per pound of body weight per day, depending on severity of challenge, up to 200 mg/hd/day monensin.

CATTLE WEIGHT (pounds)	LBS. TO FEED PER DAY MINIMUM	LBS. TO FEED PER DAY MAXIMUM
200	0.75	2.24
300	1.12	3.36
400	1.49	4.48

CATTLE WEIGHT (pounds)	LBS. TO FEED PER DAY MINIMUM	LBS. TO FEED PER DAY MAXIMUM
500	1.87	5.33
600	2.24	5.33
700	2.61	5.33
800	2.99	5.33

PRECAUTIONARY STATEMENTS

Caution: Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning: A withdrawal period has not been established for this product in prerinuating calves. Do not use in calves to be processed for veal.

MANUFACTURED BY:

BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555

Expiration date if monensin level less than 30 g/t 10/27/2019

Net Weight on Bag and/or Invoice

RUMENSIN™ is a trademark of Eli Lilly and Company

revision 12/20/2016

CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT
 FOR INVOICE 50001 DATED 1/1/2017



**Replacement Heifer CATTLE FEED
 MEDICATED**
 SUPPLEMENT FEED CONTAINING RUMENSIN™ (MONENSIN)

FORMULA # Doe RH Dec 2016

STATEMENT OF PURPOSE

For increased rate of weight gain; for the prevention and control of coccidiosis due to Eimeria bovis and E. zuernii in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

Monensin (as monensin sodium) 75.00 grams/ton
 37.50 milligrams/pound

USE DIRECTIONS

NOTE: Monensin must fed with at least 1 pound of feed. If indicated feeding rates are less than 1 pound, mix additional non-medicated feed to make at least one (1) pound of feed.

FOR INCREASED RATE OF WEIGHT GAIN:

Feed so that cattle will be consuming between 1.33 and 5.33 pounds per head per day in not less than 1lb. of feed to provide not less than 50 nor more than 200 milligrams per head per day, or after the 5th day feed at the rate of 10.67 pounds per head per every other day in not less than 2 pounds of feed to supply 400 milligrams monensin per head every other day.

IMPORTANT: For cattle consuming monensin for the first time, during the first 5 days of feeding, cattle should receive no more than 100 mg monensin or 2.67 pounds of this feed/head/day.

FOR COCCIDIOSIS PREVENTION AND CONTROL:

Hand feed this ration daily in accordance with the feeding schedule below to supply 0.14 to 0.42 mg monensin per pound of body weight per day, depending on severity of challenge, up to 200 mg/hd/day monensin.

CATTLE WEIGHT (pounds)	LBS. TO FEED PER DAY MINIMUM	LBS. TO FEED PER DAY MAXIMUM
200	0.75	2.24
300	1.12	3.36
400	1.49	4.48

CATTLE WEIGHT (pounds)	LBS. TO FEED PER DAY MINIMUM	LBS. TO FEED PER DAY MAXIMUM
500	1.87	5.33
600	2.24	5.33
700	2.61	5.33
800	2.99	5.33

PRECAUTIONARY STATEMENTS

Caution: Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning: A withdrawal period has not been established for this product in prerinuating calves. Do not use in calves to be processed for veal.

MANUFACTURED BY:

BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555

Expiration date if monensin level less than 30 g/t 10/27/2019

Net Weight on Bag and/or Invoice

RUMENSIN™ is a trademark of Eli Lilly and Company

revision 12/20/2016