

CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT

FOR INVOICE 16565 DATED 01/01/17

Green Meadows Calf Starter Cattle Feed
Medicated Feed Containing Monensin and the Feed Additive Diflubenzuron

FORMULA GMCS Sept 16



STATEMENT OF PURPOSE

For increased rate of weight gain and for prevention and control of coccidiosis due to Eimeria bovis and E. zuernii in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

For the prevention of adult house, stable, face and horn fly, emergence from the manure of treated cattle.

ACTIVE INGREDIENTS

Monensin	35.00 grams/ton	or	17.50 mg/pound
Diflubenzuron	3.35 grams/ton	or	1.67 mg/pound

USE DIRECTIONS

Feed to supply diflubenzuron at a rate of 0.1 mg/2.2 lbs body wt/day as indicated in the chart below. If the monensin chart to the right is blank, the feed cannot be used for animals of that weight; it will over or under medicate.

For growing cattle on pasture or in dry lot, according to the chart below, feed to provide 0.14 - 0.42 mg/lb. of bodyweight per day up to 200 mg/head/day depending on the severity of challenge. During the first 5 days of feeding, cattle should receive no more than 100 mg per day contained in not less than 1 lb. of feed.

Cattle Weight (pounds)	Diflubenzuron required (mg)	lbs. to feed per day
100	4.55	2.71
200	9.09	5.43
300	13.64	8.14
400	18.18	10.85
500	22.73	13.57
600	27.27	16.28
700	31.82	19.00
800	36.36	21.71
900	40.91	24.42
1000	45.45	27.14
1100	50.00	29.85
1200	54.55	32.56

Pasture/Dry Lot				
Approved level required (mg)	Monensin level in feed			
50-200				
50-200	94.98			
50-200	142.47			
50-200	189.96			
50-200				
50-200				
50-200				
50-200				
50-200				
50-200				
50-200				
50-200				
50-200				
50-200				
50-200				
50-200				

PRECAUTIONARY STATEMENTS

Caution: Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION-KEEP OUT OF REACH OF CHILDREN

See the attached diflubenzuron label related to safety (i.e. personal protective equip., first aid environmental hazards etc.).

Made by: BLUE BIRD MILL, ANYTOWN, ANY STATE

See invoice for quantity/weight

Last Revision 12/19/2016

CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT

FOR INVOICE 16565 DATED 01/01/17

Green Meadows Calf Starter Cattle Feed
Medicated Feed Containing Monensin and the Feed Additive Diflubenzuron



FORMULA GMCS Sept 16

STATEMENT OF PURPOSE

For increased rate of weight gain and for prevention and control of coccidiosis due to Eimeria bovis and E. zuernii in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).
 For the prevention of adult house, stable, face and horn fly, emergence from the manure of treated cattle.

ACTIVE INGREDIENTS

Monensin	35.00 grams/ton	or	17.50 mg/pound
Diflubenzuron	3.35 grams/ton	or	1.67 mg/pound

USE DIRECTIONS

Feed to supply diflubenzuron at a rate of 0.1 mg/2.2 lbs body wt/day as indicated in the chart below. If the monensin chart to the right is blank, the feed cannot be used for animals of that weight; it will over or under medicate.

For growing cattle on pasture or in dry lot, according to the chart below, feed to provide 0.14 - 0.42 mg/lb. of bodyweight per day up to 200 mg/head/day depending on the severity of challenge. During the first 5 days of feeding, cattle should receive no more than 100 mg per day contained in not less than 1 lb. of feed.

Cattle Weight (pounds)	Diflubenzuron required (mg)	lbs. to feed per day	Pasture/Dry Lot	
			Approved level required (mg)	Monensin level in feed
100	4.55	2.71	50-200	
200	9.09	5.43	50-200	94.98
300	13.64	8.14	50-200	142.47
400	18.18	10.85	50-200	189.96
500	22.73	13.57	50-200	
600	27.27	16.28	50-200	
700	31.82	19.00	50-200	
800	36.36	21.71	50-200	
900	40.91	24.42	50-200	
1000	45.45	27.14	50-200	
1100	50.00	29.85	50-200	
1200	54.55	32.56	50-200	

PRECAUTIONARY STATEMENTS

Caution: Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION-KEEP OUT OF REACH OF CHILDREN

See the attached label related to safety (i.e. personal protective equip., first aid environmental hazards etc.).

Made by: BLUE BIRD MILL, ANYTOWN, ANY STATE

See invoice for quantity/weight

Last Revision 12/19/2016