



TAKE TIME		CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT		TAKE TIME	
		FOR INVOICE	160091	DATED	01/01/17
OBSERVE LABEL DIRECTIONS					
<b>Jones Beef Finisher CATTLE FEED</b>					
<b>MEDICATED</b>					
COMPLETE FEED CONTAINING THE DRUGS RUMENSIN™(monensin) AND TYLAN™(tylosin)					
<b>CAUTION:</b> Federal law restricts medicated feed containing this VFD drug to use by or on the order of a licensed veterinarian.					
<b>VFD Expiration Date: Saturday, January 7, 2017</b>					
FORMULA # JB 04-15-2016					
<b>STATEMENT OF PURPOSE</b>					
For improved feed efficiency and for reduction of incidence of liver abscesses caused by <i>Fusobacterium necrophorum</i> and <i>Arcanobacterium (Actinomyces) pyogenes</i> .					
<b>ACTIVE DRUG INGREDIENTS</b>					
Monensin (monensin sodium)		5.09 grams/ton			
		2.54 mg/pound			
Tylosin phosphate		8.98 grams/ton			
		4.49 mg/pound			
<b>USE DIRECTIONS</b>					
Feed continuously as sole ration so that each animal is consuming from <b>13.36</b>					
to <b>20.04</b> pounds per head per day which will supply 60 to 90 milligrams					
tylosin and <b>34.00</b> to <b>51.00</b> milligrams monensin per head per day.					
No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 grams per ton (360 milligrams per head per day).					
<b>NOTICE: This feed may not be fed after Saturday, January 7, 2017</b>					
<b>PRECAUTIONARY STATEMENTS</b>					
<b>Caution:</b> Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.					
<b>Warning:</b> A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.					
MANUFACTURED BY:					
BLUE BIRD FEED MILL, ANY CITY ANY STATE					
Net Weight on Bag and/or Invoice					
Rumensin™ and Tylan™ are registered trademarks of Eli Lilly Company. <span style="float: right;">3/2/2017</span>					

TAKE TIME		CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT		TAKE TIME	
		FOR INVOICE	160091	DATED	01/01/17
OBSERVE LABEL DIRECTIONS					
<b>Jones Beef Finisher CATTLE FEED</b>					
<b>MEDICATED</b>					
COMPLETE FEED CONTAINING THE DRUGS RUMENSIN™(monensin) AND TYLAN™(tylosin)					
<b>CAUTION:</b> Federal law restricts medicated feed containing this VFD drug to use by or on the order of a licensed veterinarian.					
<b>VFD Expiration Date: Saturday, January 7, 2017</b>					
FORMULA # JB 04-15-2016					
<b>STATEMENT OF PURPOSE</b>					
For improved feed efficiency and for reduction of incidence of liver abscesses caused by <i>Fusobacterium necrophorum</i> and <i>Arcanobacterium (Actinomyces) pyogenes</i> .					
<b>ACTIVE DRUG INGREDIENTS</b>					
Monensin (monensin sodium)		5.09 grams/ton			
		2.54 mg/pound			
Tylosin phosphate		8.98 grams/ton			
		4.49 mg/pound			
<b>USE DIRECTIONS</b>					
Feed continuously as sole ration so that each animal is consuming from <b>13.36</b>					
to <b>20.04</b> pounds per head per day which will supply 60 to 90 milligrams					
tylosin and <b>34.00</b> to <b>51.00</b> milligrams monensin per head per day.					
No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 grams per ton (360 milligrams per head per day).					
<b>NOTICE: This feed may not be fed after Saturday, January 7, 2017</b>					
<b>PRECAUTIONARY STATEMENTS</b>					
<b>Caution:</b> Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.					
<b>Warning:</b> A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.					
MANUFACTURED BY:					
BLUE BIRD FEED MILL, ANY CITY ANY STATE					
Net Weight on Bag and/or Invoice					
Rumensin™ and Tylan™ are registered trademarks of Eli Lilly Company. <span style="float: right;">3/2/2017</span>					