

Chapter NR 664

APPENDIX IV

COCHRAN[S APPROXIMATION TO THE BEHRENS-FISHER STUDENTS[T-TEST

Using all the available background data $c n_b$ readings, calculate the background mean $c \bar{X}_b$ and background variance $c s_b^2$. For the single monitoring well under investigation $c n_m$ reading, calculate the monitoring mean $c \bar{X}_m$ and monitoring variance $c s_m^2$.

For any set of data $c X_1, X_2, \dots, X_n$, the mean is calculated by:

$$\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n}$$

and the variance is calculated by:

$$s^2 = \frac{(X_1 - \bar{X})^2 + (X_2 - \bar{X})^2 + \dots + (X_n - \bar{X})^2}{n - 1}$$

where $X_n Y$ denotes the number of observations in the set of data.

The t-test uses these data summary measures to calculate a t-statistic $c t^*$ and a comparison t-statistic $c t_c$. The t^* value is compared to the t_c value and a conclusion reached as to whether there has been a statistically significant change in any indicator parameter.

The t-statistic for all parameters except pH and similar monitoring parameters is:

$$t^* = \frac{X_m - \bar{X}_s}{\sqrt{\frac{S_m^2}{n_m} + \frac{S_b^2}{n_b}}}$$

If the value of this t-statistic is negative then there is no significant difference between the monitoring data and background data. It should be noted that significantly small negative values may be indicative of a failure of the assumption made for test validity or errors have been made in collecting the background data.

The t-statistic $c t_c$, against which t^* will be compared, necessitates finding t_b and t_m from standard cone-tailed tables where,

t_b =t-tables with $c n_b - 1$ d degrees of freedom, at the 0.05 level of significance.

t_m =t-tables with $c n_m - 1$ d degrees of freedom, at the 0.05 level of significance.

Finally, the special weightings W_b and W_m are defined as:

$$W_b = \frac{S_b^2}{n_b} \quad \text{and} \quad W_m = \frac{S_m^2}{n_m}$$

and so the comparison t-statistic is:

$$t_c = \frac{W_b t_b + W_m t_m}{W_b + W_m}$$

The t-statistic $c t^*$ is now compared with the comparison t-statistic $c t_c$ using the following decision-rule:

If t^* is equal to or larger than t_c , then conclude that there most likely has been a significant increase in this specific parameter.

If t^* is less than t_c , then conclude that most likely there has not been a change in this specific parameter.

The t-statistic for testing pH and similar monitoring parameters is constructed in the same manner as previously described except the negative sign if any is discarded and the caveat concerning the negative value is ignored. The standard c 2-tailed tables are used in the construction t_c for pH and similar monitoring parameters.

If t^* is equal to or larger than t_c , then conclude that there most likely has been a significant increase if the initial t^* had been negative, this would imply a significant decrease. If t^* is less than t_c , then conclude that there most likely has been no change.

A further discussion of the test may be found in *Statistical Methods* c6th Edition, Section 4.14d by G. W. Snedecor and W. G. Cochran, or *Principles and Procedures of Statistics* c1st Edition, Section 5.8d by R. G. D. Steel and J. H. Torrie.

STANDARD T—TABLES 0.05 LEVEL OF SIGNIFICANCE

Degrees of freedom	t-values cone-taild	t-values c 2-taild
1.....	6.314	12.706
2.....	2.920	4.303
3.....	2.353	3.182
4.....	2.132	2.776
5.....	2.015	2.571
6.....	1.943	2.447
7.....	1.895	2.365
8.....	1.860	2.306
9.....	1.833	2.262
10.....	1.812	2.228
11.....	1.796	2.201
12.....	1.782	2.179
13.....	1.771	2.160
14.....	1.761	2.145
15.....	1.753	2.131
16.....	1.746	2.120
17.....	1.740	2.110
18.....	1.734	2.101
19.....	1.729	2.093
20.....	1.725	2.086
21.....	1.721	2.080
22.....	1.717	2.074
23.....	1.714	2.069
24.....	1.711	2.064
25.....	1.708	2.060
30.....	1.697	2.042
40.....	1.684	2.021

Adopted from Table III of XStatistical Tables for Biological, Agricultural, and Medical ResearchY c1947, R. A. Fisher and F. Yatesd.